

**AMENDMENTS TO THE CLAIMS**

The following listing of claims replaces all prior versions of claims in the application.

1. (Currently Amended): A solvent for dissolving nucleic acids, ~~acid, characterized by~~  
comprising:

an ionic liquid which can dissolve nucleic ~~acid~~ acids, said ionic liquid comprising:

at least one cation selected from the group consisting of imidazolium cations and  
pyridinium cations, and

an anion which is selected from the group consisting of  $\text{BF}_4^-$ ,  $\text{PF}_6^-$ ,  $\text{AsF}_6^-$ ,  $\text{SbF}_6^-$ ,  $\text{AlCl}_4^-$ ,  
 $\text{HSO}_4^-$ ,  $\text{ClO}_4^-$ ,  $\text{CH}_3\text{SO}_3^-$ ,  $\text{CF}_3\text{SO}_3^-$ ,  $(\text{CF}_3\text{SO}_2)_2\text{N}^-$ , a halide ion and a carboxylic acid ion having a  
total of 1 to 3 carbons.

2. (Cancelled)

3. (Currently Amended): The solvent for dissolving nucleic ~~acid~~ acids of claim 1 or 10, wherein  
~~2 which is characterized in that the ionic liquid is composed of an~~ said anion which is selected  
from the group consisting of said [[a]] halide ion [[or a]] and said carboxylic acid ion having a  
total of 1 to 3 carbons.

4. (Currently Amended): The solvent for dissolving nucleic ~~acid~~ acids of claim 1 or 10, wherein  
~~which is characterized in that the ionic liquid is a neutralized ionic liquid.~~

5. (Currently Amended): The solvent for dissolving nucleic ~~acid~~ acids of claim 1 or 10, wherein  
~~or 2 which is characterized by being~~ said solvent is adapted to preserve nucleic ~~acid~~ acids or to  
react nucleic ~~acid~~ acids.
6. (Currently Amended): A nucleic acid-containing solution, ~~characterized by~~ comprising  
nucleic ~~acid~~ acids dissolved in an ionic liquid.
7. (Currently Amended): A method for preserving nucleic ~~acid~~ acids, ~~characterized by~~  
comprising the step of preserving nucleic ~~acid~~ acids in a dissolved state within an ionic liquid for  
a long term.
8. (New): The method for preserving nucleic acids of claim 7, wherein said long term is at least  
48 hours.
9. (New): The method for preserving nucleic acids of claim 7, wherein said long term is at least  
120 hours.
10. (New): The solvent for dissolving nucleic acids of claim 1, wherein said anion is selected  
from the group consisting of  $\text{PF}_6^-$ ,  $\text{AsF}_6^-$ ,  $\text{SbF}_6^-$ ,  $\text{AlCl}_4^-$ ,  $\text{HSO}_4^-$ ,  $\text{ClO}_4^-$ ,  $\text{CH}_3\text{SO}_3^-$ ,  $\text{CF}_3\text{SO}_3^-$ ,  
 $(\text{CF}_3\text{SO}_2)_2\text{N}^-$ , said halide ion and said carboxylic acid ion having a total of 1 to 3 carbons.

11. (New) A method of dissolving nucleic acids, comprising the step of:

dissolving nucleic acids with an ionic liquid which can dissolve nucleic acids,

wherein said ion liquid comprises:

at least one cation selected from the group consisting of imidazolium cations and pyridinium cations, and

an anion which is selected from the group consisting of  $\text{BF}_4^-$ ,  $\text{PF}_6^-$ ,  $\text{AsF}_6^-$ ,  $\text{SbF}_6^-$ ,  $\text{AlCl}_4^-$ ,  $\text{HSO}_4^-$ ,  $\text{ClO}_4^-$ ,  $\text{CH}_3\text{SO}_3^-$ ,  $\text{CF}_3\text{SO}_3^-$ ,  $(\text{CF}_3\text{SO}_2)_2\text{N}^-$ , a halide ion and a carboxylic acid ion having a total of 1 to 3 carbons.

12. (New): The method of dissolving nucleic acids of claim 11, wherein said anion is selected from the group consisting of  $\text{PF}_6^-$ ,  $\text{AsF}_6^-$ ,  $\text{SbF}_6^-$ ,  $\text{AlCl}_4^-$ ,  $\text{HSO}_4^-$ ,  $\text{ClO}_4^-$ ,  $\text{CH}_3\text{SO}_3^-$ ,  $\text{CF}_3\text{SO}_3^-$ ,  $(\text{CF}_3\text{SO}_2)_2\text{N}^-$ , said halide ion and said carboxylic acid ion having a total of 1 to 3 carbons.

13. (New): The method of dissolving nucleic acids of claim 11 or 12, wherein said anion is selected from the group consisting of said halide ion and said carboxylic acid ion having a total of 1 to 3 carbons.

14. (New): The method of dissolving nucleic acids of claim 11 or 12, wherein the ionic liquid is a neutralized ionic liquid.